

Immunizations	Age									
	Birth	1 month	2 months	4 months	6 months	9 months	12 months	15 months	18 months	
Hepatitis B (Hep B)	•		•	•	•					
Rotavirus			•	•	•					
Haemophilus influenzae type B (Hib)			•	•				•		
Pneumococcal conjugate (PCV)			•	•	•			•		
Diphtheria/tetanus/pertussis (DTaP)			•	•	•			•		
Inactivated polio virus (IPV)			•	•	•					
COVID-19					COVID-19*					
Measles/mumps/rubella (MMR)							•			
Varicella (chickenpox)							•			
Hepatitis A (Hep A)							•		•	
Influenza (during flu season)					Annual vaccine, 1 or 2 doses					
Age										
	4 years	5 years	6 years	10 years	11 years	12 years	16 years	17 years	18 years	
Hepatitis B (Hep B)										
Rotavirus										
Haemophilus influenzae type B (Hib)										
Pneumococcal conjugate (PCV)										
Tetanus/diphtheria/pertussis (Tdap)	•				•					
Inactivated polio virus (IPV)	•									
COVID-19	COVID-19*									
Measles/mumps/rubella (MMR)	•									
Varicella (chickenpox)	•									
Hepatitis A (Hep A)										
Human papillomavirus (HPV)					2 to 3 doses, depending on patient's age					
Meningococcal conjugate (MCV)					•		•			
Meningococcal B (MenB)							2 doses, situation-dependent			
Influenza (during flu season)	Annual vaccine, 1 or 2 doses				Annual vaccine, 1 dose starting at 8 years old					

*Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

† Influenza: Children 6 months through 8 years old who get influenza vaccine for the first time should get two doses—the second dose should be given at least four weeks after the first dose.